## Total number of printed pages-8

3 (Sem-4/CBCS) MAT HC3

2024

## MATHEMATICS

(Honours Core)

Paper: MAT-HC-4036

(Ring Theory)

Full Marks: 80

Time: Three hours

## The figures in the margin indicate full marks for the questions.

- 1. Answer the following questions as directed:  $1 \times 10 = 10$ 
  - (a) Give an example to show that for two non-zero elements a and b of a ring R, the equation ax = b can have more than one solution.
  - (b) How many nilpotent elements have in an integral domain?

- (c) Which of the following statements is not true?
  - (i)  $\langle 5 \rangle$  is a prime ideal of Z.
  - (ii)  $\langle 5 \rangle$  is a maximal ideal of Z.
  - (iii)  $\langle 5 \rangle$  is a maximal ideal of  $Z_{20}$ .
  - (iv)  $\frac{Z}{5Z}$  is an integral domain.
- (d) Define prime ideal of a ring.
- (e) Give example of a commutative ring without zero divisor that is not an integral domain.
- (f) Consider the polynomial  $f(x) = 4x^3 + 2x^2 + x + 4 \text{ and}$   $g(x) = 3x^4 + 3x^3 + 3x^2 + x + 4 \text{ in } Z_5.$  Compute f(x) + g(x).
- (g) Write  $f(x) = x^3 + x^2 + x + 1 \in \mathbb{Z}_2[x]$  as a product of irreducible polynomial over  $\mathbb{Z}_2$ .

(h) Which of the following is a primitive polynomial?

(i) 
$$2x^3 + 4x^2 + 6x + 10$$

(ii) 
$$5x^2 - 30x - 20$$

(iii) 
$$2x^4 + 3x^3 + 5x^2 - 7x + 11$$

(iv) 
$$3x^2 - 3x + 3$$

- (i) State whether the following statement is true or false:
  - "A polynomial f(x) in Z[x] which is reducible over Z is also reducible over O."
- (i) Choose the correct statement:
  - (i) Every Euclidean domain is a unique factorization domain.
  - (ii) Every principal ideal domain is a Euclidean domain.
  - (iii) Every unique factorization domain is a Euclidean domain.
  - (iv) Every unique factorization domain is a principal ideal domain.

3

- 2. Answer the following questions:  $2 \times 5 = 10$ 
  - (a) If a and b are two idempotents in a commutative ring, then show that a+b-ab is also an idempotent element.
  - (b) Show that every non-zero element of  $Z_n$  is a unit or a zero divisor.
  - (c) Show that every ring homomorphism  $f: Z_n \to Z_n$  is of the form f(x) = ax where  $a = a^2$ .
  - (d) Find the zeros of  $f(x) = x^2 + 3x + 2$  in  $Z_6$ .
  - (e) Let D be an integral domain and  $a,b \in D$ . If  $\langle a \rangle = \langle b \rangle$ , then show that a and b are associates.
- 3. Answer **any four** questions:  $5\times 4=20$ 
  - (a) The operations  $\oplus$  and  $\otimes$  defined on the set Z of integers by  $a \oplus b = a + b 1$  and  $a \otimes b = a + b ab$ . Show that  $(Z, \oplus, \otimes)$  is a ring with unity.
  - (b) Find all ring homomorphism from  $Z \oplus Z$  to Z.

- (c) Let R be a commutative ring with unity. Show that an ideal A of R is prime if and only if the quotient ring  $\frac{R}{A}$  is an integral domain.
- (d) Define principal ideal domain. Show that if F is a field, then F[x] is a principal ideal domain. 1+4=5
- (e) Show that every Euclidean domain is a principal ideal domain.
- (f) Show that the number of reducible polynomials over  $Z_p$  of the form  $x^2 + ax + b$  is  $\frac{p(p+1)}{2}$ .
- 4. Answer the following questions: 10×4=40

5

(a) (i) Let R be a commutative ring with unity. Show that the set

$$R[x] = \{a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 / a_i \in R ,$$

n is a non-negative integer} is a ring. Also show that if R is an integral domain, then R[x] is also an integral domain. 5+2=7

(ii) Let 
$$f(x) = 5x^4 + 3x^3 + 1$$
 and  $g(x) = 3x^2 + 2x + 1$  in  $Z_7[x]$ .

Determine the quotient and remainder upon dividing f(x) by g(x).

Or

(i) Show that

$$Z\left|\sqrt{2}\right| = \left\{a + b\sqrt{2} : a, b \in Z\right\}$$
 and  $H = \left\{\begin{bmatrix} a & 2b \\ b & a \end{bmatrix} / a, b \in Z\right\}$ 

are isomorphic as ring.

4

- (ii) If a, b be any two ring elements and m and n be any two integers, then show that (m.a)(n.b) = (mn).(ab)
- (b) (i) Define maximal ideal of a ring. Let A be an ideal of a commutative ring with unity R. Prove that  $\frac{R}{A}$  is a field if and only if A is maximal. 1+6=7

(ii) Let R be a commutative ring and A be any subset of R. Show that the nil-radical of A,

 $N(A) = \{ r \in R / r^n \in A \text{ for some } n \in N \}$  is an ideal of R.

Or

- (i) Let  $\varphi$  be a ring homomorphism from R to S. Then the mapping  $\frac{R}{\ker(\varphi)} \text{ to } \varphi(R), \text{ given by } r + \ker(\varphi) \to \varphi(r) \text{ is an }$ isomorphism, i.e.,  $\frac{R}{\ker(\varphi)} \approx \varphi(R)$ .
- (ii) Let  $\phi$  be a ring homomorphism from a ring R to a ring S. Let B be an ideal of S. Then  $\phi^{-1}[B] = \{r \in R : \phi(r) \in B\}$  is an ideal of R.
- (c) If F is a field and  $p(x) \in F[x]$ , then prove that  $\frac{F[x]}{\langle p(x) \rangle}$  is a field if and only if p(x) is irreducible over F.

Let F be a field. If  $f(x) \in F[x]$  and degree f(x) is 2 or 3, then f(x) is reducible over F if and only if f(x) has a zero in F. Is the result true when degree f(x) is greater then 3? Justify.

7+3=10

(d) In a principal ideal domain, show that an element is irreducible if and only if it is prime. Use this result to show that  $Z\left[\sqrt{-3}\right] = \left\{a + b\sqrt{-3} : a, b \in Z\right\}$  is not a principal ideal domain. 7+3=10

## Or

- (i) In a principal ideal domain, show that any strictly increasing chain of ideals  $I_1 \subset I_2 \subset ...$  must be finite in length.
- (ii) Let  $\phi$  be a onto ring homomorphism from a ring R to a ring S. Then prove that  $\phi$  is an isomorphism if and only if  $\ker(\phi) = \{0\}$ .
- (iii) Determine all ring homomorphism from the ring of integers Z to itself.