

Total number of printed pages-7

3 (Sem-5/CBCS) MAT HC 1

2024

MATHEMATICS

(Honours Core)

(New Course)

Paper : MAT-HC-5016

(Complex Analysis)

Full Marks : 60

Time : Three hours

***The figures in the margin indicate
full marks for the questions.***

1. Answer the following questions : $1 \times 7 = 7$

(a) The function

$$f(z) = xy^2 + e^{xy} + i(2x - y) \text{ is}$$

continuous everywhere in the complex plane.

(State True or False)

Contd.

1-2-2023

(b) Let the function $f(z) = u(x, y) + iv(x, y)$ be defined throughout some ε -neighbourhood of a point $z_0 = x_0 + iy_0$. State a sufficient condition for existence of the derivative $f'(z_0)$.

(c) Find $\lim_{z \rightarrow -1} \frac{iz + 3}{z + 1}$.

(d) Define entire function and give an example.

(e) Show that $\exp(2 \pm 3\pi i) = -e^2$.

(f) What is a Jordan curve?

(g) State Liouville's theorem.

2. Answer the following questions : $2 \times 4 = 8$

(a) Using ε - δ definition, show that if

$$f(z) = z^2 \text{ then } \lim_{z \rightarrow z_0} f(z) = z_0^2.$$

(b) Show that $f(z) = \begin{cases} z^2, & z \neq z_0 \\ 0, & z = z_0 \end{cases}$ where

$z_0 \neq 0$ is discontinuous at $z = z_0$.

(c) Determine the singular points of the function, $f(z) = \frac{2z+1}{z(z^2+1)}$.

(d) Evaluate $\int_C \bar{z} dz$ from $z=0$ to $z=4+2i$

along the curve C given by $z = t^2 + it$.

3. Answer **any three** questions from the following : $5 \times 3 = 15$

(a) Show that the three cube roots of $-8i$ lie at the vertices of an equilateral triangle that is inscribed in a circle of radius 2 centred at the origin.

(b) Prove that $\lim_{z \rightarrow 0} \frac{\bar{z}}{z}$ does not exist.

(c) If $f'(z) = 0$ everywhere in a domain D , then prove that $f(z)$ must be constant throughout D .

(d) Suppose that a function $f(z)$ is analytic at a point $z_0 = z(t_0)$ on a differentiable arc $z = z(t) (a \leq t \leq b)$. Show that if $w(t) = f(z(t))$ then $w'(t) = f'(z(t))z'(t)$ when $t = t_0$.

(e) Using anti-derivative, evaluate the integral $\int_C z^{1/2} dz$, where C is a contour from $z = -3$ to $z = 3$ that, except for its end points, lies above the x -axis.

4. Answer **any three** questions from the following : 10×3=30

(a) (i) If z_0 and w_0 are points in the z and w planes respectively, then prove that

(A) $\lim_{z \rightarrow z_0} f(z) = \infty$ if and only if

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0 \text{ and}$$

(B) $\lim_{z \rightarrow \infty} f(z) = w_0$ if and only if

$$\lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = w_0 \quad 5$$

(ii) Let u and v denote the real and imaginary components of the function f defined by the equations :

$$f(z) = \begin{cases} \frac{(\bar{z})^2}{z}, & \text{when } z \neq 0 \\ 0, & \text{when } z = 0 \end{cases}$$

Verify the Cauchy-Riemann equations at the origin $z = (0,0)$.

5

(b) (i) Show that

$$\sin(x+iy) = \sin x \cosh y + i \cos x \sinh y.$$

2

(ii) Find numbers $z = x+iy$ such that $e^z = 1+i$.

3

(iii) Show that if a function

$$f(z) = u(x,y) + iv(x,y)$$

and its conjugate $\overline{f(z)}$ are both analytic in a domain D , then $f(z)$ must be constant throughout D .

5

(c) (i) Show that the zeros of $\sin z$ are all real.

2

(ii) Evaluate $\int_0^{7/4} e^t dt$.

3

(iii) If $w = f(z) = \frac{1+z}{1-z}$ find $\frac{dw}{dz}$ and determine where $f(z)$ is not analytic.

5

- (d) (i) If $w(t)$ is a piecewise continuous complex valued function defined on an interval $a \leq t \leq b$, then show

$$\text{that } \left| \int_a^b w(t) dt \right| \leq \int_a^b |w(t)| dt \quad 5$$

- (ii) Let C denote the line segment from $z = i$ to $z = 1$. By observing that of all the points on that line segment, the midpoint is the closest to the

origin, show that $\left| \int_C \frac{dz}{z^4} \right| \leq 4\sqrt{2},$

without evaluating the integral.

- (e) (i) Show that

$$\int \frac{dz}{z^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{z}{a} + C_1 = \frac{1}{2ai} \ln \left(\frac{z - ai}{z + ai} \right) + C_2$$

- (ii) Evaluate :

$$\oint_C \frac{\sin \pi z^2 + \cos \pi z^2}{(z-1)(z-2)} dz, \quad 5$$

where C is the circle $|z| = 3$.

- (f) (i) Prove that if a function f is analytic at a given point, then its derivatives of all orders are also analytic at that point. 5

- (ii) Let C denote the positively oriented boundary of a square whose sides lie along the lines $x = \pm 2$ and $y = \pm 2$. Evaluate

$$\int_C \frac{\tan(z/2)}{(z-x_0)^2} dz \quad (-2 < x_0 < 2). \quad 5$$