

2 0 1 8

PHYSICS

(Major)

Paper : 5.2

(Atomic Physics)

Full Marks : 60

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. Choose the correct option of any *seven* of the following : 1×7=7

(a) An electron revolves about a proton in second excited state. The angular momentum of the electron is

(i) $\frac{h}{2\pi}$

(ii) $\frac{h}{\pi}$

(iii) $\frac{3h}{2\pi}$

(iv) 0

- (b) Lines of Balmer series are obtained from the hydrogen atom, when electron jumps from some higher orbit to
- (i) fourth orbit
 - (ii) third orbit
 - (iii) second orbit
 - (iv) None of the above
- (c) The formation of electronic spectrum is due to
- (i) change in electronic energy
 - (ii) change in vibrational energy
 - (iii) change in rotational energy
 - (iv) change in all (i), (ii) and (iii)
- (d) The minimum wavelength of X-rays produced by electrons accelerated by a potential difference of V volts is
- (i) $\frac{eV}{vc}$
 - (ii) $\frac{eV}{hc}$
 - (iii) $\frac{hc}{eV}$
 - (iv) $\frac{h}{v}$
- (e) The minimum number of electrons in a sub-shell with orbital angular momentum quantum number l is
- (i) $2(2l+1)$
 - (ii) $(2l-1)$
 - (iii) $2(2l-1)$
 - (iv) $(2l+1)$

- (f) Stern-Gerlach experiment confirms
- (i) electron spin and associated magnetic moment
 - (ii) orbital motion of the electron and associated moment
 - (iii) specific charge (e/m) of the electron
 - (iv) spin-orbit interaction of the electron
- (g) If ν_{K_α} and ν_{L_α} be the frequencies of K_α and L_α characteristic X-ray lines, then
- (i) $\nu_{K_\alpha} = \nu_{L_\alpha}$
 - (ii) $\nu_{K_\alpha} < \nu_{L_\alpha}$
 - (iii) $\nu_{K_\alpha} > \nu_{L_\alpha}$
 - (iv) $\nu_{K_\alpha} = \frac{1}{\nu_{L_\alpha}}$
- (h) The shape of the electron orbit is determined by the quantum number
- (i) n
 - (ii) l
 - (iii) j
 - (iv) m_j
- (i) The splitting of spectral lines with components in strong electric field is known as
- (i) normal Zeeman effect
 - (ii) anomalous Zeeman effect
 - (iii) Paschen-Back effect
 - (iv) Stark effect

(Turn Over)

(4)

2. Answer any *four* of the following : $2 \times 4 = 8$

- (a) A charged oil drop is suspended in an uniform electric field of 3×10^4 V/m so that it neither rises nor falls. If the mass of the drop is 9.75×10^{-15} kg, find the charge on the drop.
- (b) Find the precessional frequency of an electron orbit when placed in a magnetic field of 6 tesla. ($e = 1.6 \times 10^{-19}$ C, $m = 9.1 \times 10^{-28}$ kg)
- (c) Electron moves at right angles to magnetic field of 150×10^{-14} tesla with a velocity of 6×10^6 m/s. Find the radius of circular path. ($e/m = 1.7 \times 10^{11}$ C/kg)
- (d) What is Lande g -factor? What is the value of g -factor of an atom with a single electron in $d_{3/2}$ state.
- (e) If the PD between the anode and the cathode is 25 kV, what is the cut-off wavelength and the cut-off frequency of the emitted X-rays? ($c = 3 \times 10^8$ m/s, $h = 6.6 \times 10^{-34}$ J-s)
- (f) Using vector atom model, determine the possible values of the total angular momentum of an f -electron ($l = 3$).

(5)

3. Answer the questions (a) and any *two* from (b), (c) and (d) : $5 \times 3 = 15$

- (a) Mention the important feature of Rutherford's scattering of α -particles by gold foil which supported the nuclear model of the atom against Thomson model.
- (b) A 2 keV electron enters a magnetic field of 5×10^{-4} Wb/m². If the radius of the electron path is 0.303 m, find the (e/m) of the electron.
- (c) A beam of X-rays of wavelength 0.842 Å is incident on a crystal at a glancing angle of 8.6° , when the first-order Bragg's reflection occur. Calculate the glancing angle of the third-order reflection.
- (d) Write any *one* explanatory note on the following :
- (i) Pauli's exclusion principle
 - (ii) Alkali spectra
 - (iii) Vector atom model

4. Answer the questions (a) and (b) and any one from (c) and (d) : $10 \times 3 = 30$

- (a) If the positive charge of the gold atom is supposed to be spreaded uniformly over a spherical surface of diameter 1 \AA , show that the α -particle of energy greater than a certain value E will not be reflected back. Calculate the value of E . $\left(\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \right)$ $5+5=10$

Or

What is Compton effect? Derive an expression for the change in wavelength of a photon when it is scattered by an electron. Justify the importance of its theory. $2+5+3=10$

- (b) (i) Describe and explain L - S coupling. Under what condition does it hold?
(ii) Under what condition L - S coupling breaks down and what kind of new coupling takes place?
(iii) Describe J - J coupling. Illustrate L - S and J - J coupling with the help of vector diagram. $3+3+4=10$

Or

Using the physical constants given below, calculate the following for hydrogen atom : $2+3+3+2=10$

- (i) Velocity of an electron in the ground state
(ii) Radius of Bohr orbit in the ground state
(iii) Time taken by the electron to transverse first orbit
(iv) Rydberg constant

$$C = 1.6 \times 10^{-19} \text{ C}, \quad m = 9.1 \times 10^{-31} \text{ kg}$$

$$h = 6.6 \times 10^{-34} \text{ J-s}, \quad c = 3 \times 10^8 \text{ m/s}$$

$$\epsilon_0 = 8.86 \times 10^{-12} \text{ C}^2 / \text{N-m}^2$$

- (c) (i) Enumerate briefly the theory of Raman effect.
(ii) Why are the Stokes lines brighter than the anti-Stokes lines?
(iii) Compare Raman spectra with infrared spectra. $4+3+3=10$
- (d) What is Zeeman effect? Draw a neat diagram to illustrate the Zeeman splitting of D_1 and D_2 lines of sodium. What is the difference between normal and anomalous Zeeman effect? Write the Zeeman shift in terms of wavelength and e/m in terms of Zeeman shift. $2+3+3+2=10$

★ ★ ★