

Total number of printed pages-8

3 (Sem-5/CBCS) MAT HC 1

2021

(Held in 2022)

MATHEMATICS

(Honours)

Paper : MAT-HC-5016

(Riemann Integration and Metric Spaces)

Full Marks : 80

Time : Three hours

**The figures in the margin indicate
full marks for the questions.**

1. Answer the following as directed : $1 \times 10 = 10$

(a) Describe an open ball in the discrete metric space.

(b) Find the derived set of the sets $(0, 1]$ and $[0, 1]$.

(c) A subset B of a metric space (X, d) is open if and only if

(i) $B = \overline{B}$

(ii) $B = B^\circ$

(iii) $B \neq \overline{B}$

(iv) $B \neq B^\circ$

(Choose the correct one)

Contd.

Lib. Set
5016, 5026.

(d) Which of the following is false ?

- (i) $\phi^o = \phi, X^o = X$
- (ii) $A \subseteq B \Rightarrow A^o \subseteq B^o$
- (iii) $(A \cap B)^o = A^o \cap B^o$
- (iv) $(A \cup B)^o = A^o \cup B^o$

where A, B are subsets of a metric space (X, d) . (Choose the false one)

(e) The closure of the subset

$F = \left\{1, \frac{1}{2}, \frac{1}{3}, \dots\right\}$ of the real line \mathbb{R} is

- (i) ϕ
- (ii) F
- (iii) $F \cup \{0\}$
- (iv) $F - \{0\}$

(Choose the correct one)

(f) In a metric space an arbitrary union of closed sets need not be closed. Justify it with an example.

(g) If A is a subset of a metric space (X, d) , then which one is true ?

- (i) $d(A) = d(\overline{A})$
- (ii) $d(A) \neq d(\overline{A})$
- (iii) $d(A) > d(\overline{A})$
- (iv) $d(A) < d(\overline{A})$

(Choose the true one)

(h) When is an improper Riemann integral said to be convergent ?

- (i) Evaluate $\int_0^{\infty} e^{-x} dx$ if it exists.
- (j) Show that $\Gamma(1) = 1$

2. Answer the following questions : $2 \times 5 = 10$

(a) Let F be a subset of a metric space (X, d) . Prove that the set of limit points of F is a closed subset of (X, d) .

(b) If F_1 and F_2 are two subsets of a metric space (X, d) , then

$\overline{F_1 \cap F_2} = \overline{F_1} \cap \overline{F_2}$. Justify whether it is false or true.

(c) Let (X, d_X) and (Y, d_Y) be metric spaces and $f: X \rightarrow Y$. If for all subsets A of X , $f(\overline{A}) \subseteq \overline{f(A)}$, then show that f is continuous on X .

(d) Let $f: [a, b] \rightarrow \mathbb{R}$ be integrable. Show that $|f|$ is integrable.

(e) Show that the function $f: [a, b] \rightarrow \mathbb{R}$ defined by $f(x) = c$ for all $x \in [a, b]$ is integrable with its integral $c(b-a)$.

3. Answer **any four** parts : 5×4=20

(a) Define a complete metric space. Show that the metric space $X = \mathbb{R}^n$ with the metric given by

$$d_p(x, y) = \left(\sum |x_i - y_i|^p \right)^{\frac{1}{p}}, \quad p \geq 1$$

where $x = (x_1, x_2, \dots, x_n)$ and

$y = (y_1, y_2, \dots, y_n)$ are in \mathbb{R}^n , is a complete metric space. 1+4=5

(b) Let (X, d_X) and (Y, d_Y) be metric spaces. Prove that a mapping $f: X \rightarrow Y$ is continuous on X if and only if $f^{-1}(G)$ is open in X for all open subsets G of Y . 5

(c) Prove that if the metric space (X, d) is disconnected, then there exists a continuous mapping of (X, d) onto the discrete two-element space (X_0, d_0) . 5

(d) Let $f: [a, b] \rightarrow \mathbb{R}$ be a continuous function. Prove that f is integrable. 5

(e) Discuss the convergence of the integral $\int_1^{\infty} \frac{1}{x^p} dx$ for various values of p . 5

(f) Show that for $a > -1$,

$$S_n = \frac{1^n + 2^n + \dots + n^n}{n^{1+a}} \rightarrow \frac{1}{1+a}. \quad 5$$

4. Answer **any four** parts : $10 \times 4 = 40$

(a) (i) Let (X, d) be a metric space. Define $d : X \times X \rightarrow \mathbb{R}$ by

$$d'(x, y) = \frac{d(x, y)}{1 + d(x, y)} \text{ for all}$$

$x, y \in X$. Prove that d' is a metric on X .

Also show that d and d' are equivalent metrics on X .

4+2=6

(ii) Prove that a convergent sequence in a metric space is a Cauchy sequence.

4

(b) (i) Let (X, d) be a metric space and F be a subset of X . Prove that F is closed in X if and only if F^c is open.

5

(ii) If (Y, d_Y) is a subspace of a metric space (X, d) , then show that a subset Z of Y is open in Y if and only if there exists an open set $G \subseteq X$ such that $Z = G \cap Y$.

5

(c) Prove that a metric space (X, d) is complete if and only if for every nested sequence $\{F_n\}_{n \geq 1}$ of non-empty closed subsets of X such that $d(F_n) \rightarrow 0$ as

$n \rightarrow \infty$, the intersection $\bigcap_{n=1}^{\infty} F_n$ contains one and only one point.

10

(d) (i) Prove that in a metric space (X, d) , each open ball is an open set.

4

(ii) Let (X, d_X) and (Y, d_Y) be metric spaces and $A \subseteq X$. Prove that a function $f : A \rightarrow Y$ is continuous at $a \in A$ if and only if whenever a sequence $\{x_n\}$ in A converges to a , the sequence $\{f(x_n)\}$ converges to $f(a)$.

6

(e) (i) Define uniformly continuous mapping in a metric space. Give an example to show that a continuous mapping need not be uniformly continuous.

1+4=5

(ii) Prove that the image of a Cauchy sequence under a uniformly continuous mapping is itself a Cauchy sequence. 5

(f) Let (\mathbb{R}, d) be the space of real numbers with the usual metric. Prove that a subset $I \subseteq \mathbb{R}$ is connected if and only if I is an interval. 10

(g) Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Show that f is integrable if and only if it is Riemann integrable. 10

(h) (i) State and prove first fundamental theorem of calculus. Using it show that

$$\int_0^a f(x) dx = \frac{a^4}{4} \text{ for } f(x) = x^3.$$

$$1+3+2=6$$

(ii) Let f be continuous on $[a, b]$. Prove that there exists $c \in [a, b]$

$$\text{such that } \frac{1}{b-a} \int_a^b f(x) dx = f(c).$$

4

(ii) Prove that the image of a Cauchy sequence under a uniformly continuous mapping is itself a Cauchy sequence. 5

(f) Let (\mathbb{R}, d) be the space of real numbers with the usual metric. Prove that a subset $I \subseteq \mathbb{R}$ is connected if and only if I is an interval. 10

(g) Let $f: [a, b] \rightarrow \mathbb{R}$ be a bounded function. Show that f is integrable if and only if it is Riemann integrable. 10

(h) (i) State and prove first fundamental theorem of calculus. Using it show that

$$\int_0^a f(x) dx = \frac{a^4}{4} \text{ for } f(x) = x^3.$$

$$1+3+2=6$$

(ii) Let f be continuous on $[a, b]$. Prove that there exists $c \in [a, b]$

$$\text{such that } \frac{1}{b-a} \int_a^b f(x) dx = f(c).$$

4