3 (Sem-5/CBCS) MAT HC 2

2023

MATHEMATICS

(Honours Core)

Paper: MAT-HC-5026

(Linear Algebra)

Full Marks: 80

Time: Three hours

The figures in the margin indicate full marks for the questions.

 Answer the following questions as directed: 1×10=10

(a) Let
$$A = \begin{pmatrix} 1 & -3 & -2 \\ -5 & 9 & 1 \end{pmatrix}$$
 and $\vec{u} = \begin{bmatrix} 5 \\ 3 \\ -2 \end{bmatrix}$.

Check whether \vec{u} is in null space of A.

- (b) Define subspace of a vector space.
- (c) Give reason why \mathbb{R}^2 is not a subspace of \mathbb{R}^3 .

Contd.

- (d) State whether the following statement is true or false:

 "If dimension of a vector space V is p > 0 and S is a linearly dependent subset of V, then S contains more than p elements."
- (e) If \vec{x} is an eigenvector of corresponding to the eigenvalue λ then what is $A^3\vec{x}$?
- (f) When two square matrices A and Bart said to be similar?
- (g) If $\vec{v} = (1 2 \ 2 \ 4)$ then find $\|\vec{v}\|$.
- (h) Find a unit vector in the direction $\vec{u} = \begin{bmatrix} 8/3 \\ 2 \end{bmatrix}$.
- (i) Under what condition two vectors and \vec{v} are orthogonal to each other
- (i) Define orthogonal complement
- 2. Answer the following questions: 2×5^{-10}
 - (a) Show that the set $W = \left\{ \begin{bmatrix} x \\ y \end{bmatrix} : xy \ge 0 \right\}$ if not a subspace of \mathbb{R}^2 .

- (b) Let $\vec{b}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, $\vec{b}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$, $\vec{x} = \begin{bmatrix} 4 \\ 5 \end{bmatrix}$ and $\beta = \{b_1, b_2\}$. Find the coordinate vector $[x]_{\beta}$ of \vec{x} relative to β .
- (c) Find the eigenvalues of $A = \begin{bmatrix} 2 & 3 \\ 3 & -6 \end{bmatrix}$.
- (d) Let P_2 be the vector space of all polynomials of degree less than equal to 2. Consider the linear transformation $T: P_2 \to P_2$ defined by $T(a_0 + a_1t + a_2t^2) = a_1 + 2a_2t$. Find the matrix representation $[T]_\beta$ of T with respect to the base $\beta = \{1, t, t^2\}$.
- (e) Show that the matrix $A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{2}{3} \\ \frac{1}{\sqrt{2}} & -\frac{2}{3} \\ 0 & \frac{1}{3} \end{bmatrix}$

has orthogonal columns.

3

- Answer any four questions : 3. 5×4=2
 - (a) Let $S = \{v_1, v_2, ..., v_p\}$ be a set in the vector space V and H = span(S). No if one of the vector in S, say 1/1 is linear combination of the other vectors in S, then show that S linearly dependent and the subset $S_1 = S - \{v_k\}$ still span H. 2+3=
 - (b) Show that the set of all eigenvector corresponding to the disting eigenvalues of a $n \times n$ matrix A linearly independent.
 - (c) Let W be a subspace of the vector space of W Sis a linearly independent substi of W. Show that S can be extended necessary, to form a basis for Wan Answer either (a) or (b) from each of the following dim W < dim W
 - (d) If $A = \begin{bmatrix} 1 & 3 & 3 \\ -3 & -5 & -3 \\ 3 & 3 & 1 \end{bmatrix}$. Find an

invertible matrix P and a diagonal matrix D such that $A = PDP^{-1}$.

- (e) If $\vec{y} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$ and $\vec{u} = \begin{bmatrix} 4 \\ -7 \end{bmatrix}$ then find the orthogonal projection of $ar{y}$ onto $ar{u}$ and write \vec{y} as the sum of two orthogonal vectors, one in $span \{\vec{u}\}$ and the other orthogonal to \vec{u} .
 - (f) If $W = span\{x_1, x_2\}$ where $x_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$,

$$x_2 = \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \\ -\frac{2}{3} \end{bmatrix}$$
, find a orthogonal basis for W .

(a) Find a spanning set for the null space of the matrix:

$$A = \begin{bmatrix} -3 & 6 & -1 & 1 & -7 \\ 1 & -2 & 2 & 3 & -1 \\ 2 & -4 & 5 & 8 & -4 \end{bmatrix}$$

Is this spanning set linearly independent?

- (b) (i) If a vector space V has a basis n vectors, then show that even basis of V must consist of exact n vectors.
- Find a basis for column space the following matrix :

$$B = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$$

- 5. (a) Define eigenvalue and eigenvector of matrix. Find the eigenvalues corresponding eigenvectors of $\begin{array}{c|c}
 \text{matrix} & 2 & 3 \\
 3 & -6
 \end{array}$
 - (b) Let T be a linear operator on a find dimension linear operator on a find dimensional vector space V and of denote the T-cyclic subspace If dim (III) If dim(W) = k then show that
 - (i) $\{v, T(v), T^2(v), \dots, T^{k-1}(v)\}$ is basis for W.

6

- (ii) If
- $a_0v + a_1T(v) + ... + a_{k-1}T^{k-1}(v) + T^k(v) = 0$, then the characteristics polynomial of T_{m} is

$$f(t) = (-1)^k (a_0 + a_1 t + \dots + a_{k-1} t^{k-1} + t^k).$$

6+4=10

- $B = \begin{bmatrix} 1 & 4 & 0 & 2 & -1 \\ 3 & 12 & 1 & 5 & 5 \\ 2 & 8 & 1 & 3 & 2 \\ 5 & 20 & 2 & 8 & 8 \end{bmatrix}$ 6. (a) (i) Define orthogonal set of vectors. Let $S = \{\vec{u}_1, \vec{u}_2, \dots, \vec{u}_p\}$ is an arthogonal set of non-zero vectors. orthogonal set of non-zero vectors in \mathbb{R}^n , then show that S is linearly 1+4=5independent.
 - For any symmetric matrix show that (ii) any two eigenvectors from different eigenspaces are orthogonal.
 - Define inner product space. Show that the following is an inner product in \mathbb{R}^2

$$\langle u, v \rangle = 4u_1v_1 + 5u_2v_2$$

7

Where $u = (u_1, u_2), v = (v_1, v_2) \in \mathbb{R}^2$ Also, show that in any inner product space V,

$$|\langle u, v \rangle| \le ||u|| \cdot ||v||, \ \forall u, v \in V.$$

$$2+4+4=10$$

7. (a) (i) Consider the bases $\beta = \{b_1, b_1\}$ and $\gamma = \{c_1, c_2\}$ for \mathbb{R}^2 when

$$b_1 = \begin{bmatrix} 1 \\ -3 \end{bmatrix}, b_2 = \begin{bmatrix} -2 \\ 4 \end{bmatrix}, c_1 = \begin{bmatrix} -7 \\ 9 \end{bmatrix}$$

and $c_2 = \begin{bmatrix} -5 \\ 7 \end{bmatrix}$, find the charge of coordinate matrix from γ to and from β to γ .

(ii) Compute A¹⁰ where

$$A = \begin{bmatrix} 4 & -3 \\ 2 & -1 \end{bmatrix}.$$

(b) State Cayley-Hamilton theorem matrices. Verify the theorem for matrix $M = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$ and hence M^{-1} .

Total number of printed pages-8

3 (Sem-5/CBCS) MAT HC 2

2023

MATHEMATICS

(Honours Core)

Paper: MAT-HC-5026

(Linear Algebra)

Full Marks: 80

Time: Three hours

The figures in the margin indicate full marks for the questions.

Answer the following questions as directed: $1 \times 10 = 10$

(a) Let
$$A = \begin{pmatrix} 1 & -3 & -2 \\ -5 & 9 & 1 \end{pmatrix}$$
 and $\vec{u} = \begin{bmatrix} 5 \\ 3 \\ -2 \end{bmatrix}$.

Check whether \vec{u} is in null space of A.

- (b) Define subspace of a vector space.
- (c) Give reason why \mathbb{R}^2 is not a subspace of \mathbb{R}^3 .